

3-Hydroxyanilinium hydrogensulfate

Nourredine Benali-Cherif,* Alima Kateb, Houda Boussekine, Zina Boutobba and Amel Messai

Laboratoire des Structures, Propriétés et Interactions Inter Atomiques (LASPI²A), Centre Universitaire de Khenchela, 40000 Khenchela, Algeria

Correspondence e-mail: benalicherif@hotmail.com


Received 3 June 2007; accepted 13 June 2007

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.030; wR factor = 0.099; data-to-parameter ratio = 18.5.

In the title compound, $\text{C}_6\text{H}_8\text{NO}^+\cdot\text{HSO}_4^-$, there is an intricate cation–cation, cation–anion and anion–anion three-dimensional hydrogen-bond network.

Related literature

For related literature, see: Aakeroy *et al.* (1999); Benali-Cherif, Allouche *et al.* (2007); Benali-Cherif, Direm *et al.* (2007); Cherouana *et al.* (2003); Hagrman *et al.* (1999); Mazeaud *et al.* (2000); Soghomonian *et al.* (1995).



Experimental

Crystal data

$\text{C}_6\text{H}_8\text{NO}^+\cdot\text{HSO}_4^-$
 $M_r = 207.21$
 Monoclinic, $P2_1$
 $a = 7.3142$ (3) Å
 $b = 5.8612$ (2) Å
 $c = 9.8969$ (2) Å
 $\beta = 98.829$ (2)°

$V = 419.25$ (2) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 0.38$ mm⁻¹
 $T = 293$ (2) K
 $0.15 \times 0.12 \times 0.10$ mm

Data collection

Nonius KappaCCD area-detector diffractometer
 Absorption correction: none
 5851 measured reflections

2241 independent reflections
 2150 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.043$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.030$
 $wR(F^2) = 0.099$
 $S = 1.14$

2241 reflections
 121 parameters
 1 restraint

H-atom parameters constrained

$\Delta\rho_{\text{max}} = 0.32$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.45$ e Å⁻³

Absolute structure: Flack (1983),
 with 1095 Friedel pairs
 Flack parameter: -0.02 (8)

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1A}\cdots\text{O2}^{\text{ii}}$	0.89	2.03	2.830 (2)	149
$\text{N1}-\text{H1B}\cdots\text{O2}^{\text{ii}}$	0.89	1.98	2.853 (2)	166
$\text{N1}-\text{H1C}\cdots\text{O3}^{\text{iii}}$	0.89	2.14	2.962 (2)	152
$\text{N1}-\text{H1C}\cdots\text{O4}$	0.89	2.39	2.975 (2)	124
$\text{O41}-\text{H41}\cdots\text{O4}^{\text{iv}}$	0.82	1.88	2.695 (2)	174
$\text{O1}-\text{H1}\cdots\text{O41}^{\text{iv}}$	0.82	1.86	2.642 (2)	160

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z$; (ii) $x - 1, y, z$; (iii) $-x + 1, y - \frac{1}{2}, -z$; (iv) $-x + 1, y + \frac{1}{2}, -z + 1$.

Data collection: *KappaCCD Server Software* (Nonius, 1998); cell refinement: *DENZO* and *SCALEPACK* (Otwinowski & Minor, 1997); data reduction: *DENZO* and *SCALEPACK*; program(s) used to solve structure: *SIR2004* (Burla *et al.*, 2005); program(s) used to refine structure: *SHELXL97* (Sheldrick, 1997); molecular graphics: *ORTEP-3* (Farrugia, 1997) and *PLATON* (Spek, 2003); software used to prepare material for publication: *WinGX* (Farrugia, 1999).

The authors thank Dr M. Giorgi from LBS-UMR 6517, Faculté des Sciences et Techniques de Saint Jérôme, Marseille, France, for providing diffraction facilities, and the Centre Universitaire de Khenchela for financial support.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: DN2197).

References

- Aakeroy, C. B., Beatty, A. M. & Leinen, D. S. (1999). *Angew. Chem. Int. Ed.* **38**, 1815–1819.
- Benali-Cherif, N., Allouche, F., Direm, A., Boukli-H-Benmenni, L. & Soudani, K. (2007). *Acta Cryst.* **E63**, o2643–o2645.
- Benali-Cherif, N., Direm, A., Allouche, F., Boukli-H-Benmenni, L. & Soudani, K. (2007). *Acta Cryst.* **E63**, o2054–o2056.
- Burla, M. C., Caliandro, R., Camalli, M., Carrozzini, B., Cascarano, G. L., De Caro, L., Giacovazzo, C., Polidori, G. & Spagna, R. (2005). *J. Appl. Cryst.* **38**, 381–388.
- Cherouana, A., Bendjedou, L. & Benali-Cherif, N. (2003). *Acta Cryst.* **E59**, o1790–o1792.
- Farrugia, L. J. (1997). *J. Appl. Cryst.* **30**, 565.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Flack, H. D. (1983). *Acta Cryst.* **A39**, 876–881.
- Hagrman, J. P., Hagrman, D. & Zubieta, J. (1999). *Angew. Chem. Int. Ed.* **38**, 2638–2684.
- Mazeaud, A., Dromzee, Y. & Thouvenot, R. (2000). *Inorg. Chem.* **39**, 4735–4740.
- Nonius (1998). *KappaCCD Server Software*. Nonius BV, Delft, The Netherlands.
- Otwinowski, Z. & Minor, W. (1997). *Methods in Enzymology*, Vol. 276, *Macromolecular Crystallography*, Part A, edited by C. W. Carter Jr & R. M. Sweet, pp. 307–326. New York: Academic Press.
- Sheldrick, G. M. (1997). *SHELXL97*. University of Göttingen, Germany.
- Soghomonian, V., Chen, Q., Haushalter, R. C. & Zubieta, J. (1995). *Angew. Chem. Int. Ed. Engl.* **34**, 223–226.
- Spek, A. L. (2003). *J. Appl. Cryst.* **36**, 7–13.

supplementary materials

Acta Cryst. (2007). E63, o3251 [doi:10.1107/S1600536807028978]

3-Hydroxyanilinium hydrogensulfate

N. Benali-Cherif, A. Kateb, H. Boussekine, Z. Boutobba and A. Messai

Comment

Crystal engineering of organic-inorganic hybrid compounds is currently of great interest and these materials have received increasing attention during the past few decades (Mazeaud *et al.*, 2000; Soghomonian *et al.*, 1995) owing to their interesting structural topologies and potential application in materials science, such as ion-exchange, adsorption, molecular recognition, catalysis and magnetism (Aakeroy *et al.*, 1999; Hagrman *et al.*, 1999). The crystal structure of *m*-hydroxyanilinium bisulfate, (I), was determined as part of our investigations on the structural characteristics of organic-inorganic layered compounds and an ongoing study on D—H···A hydrogen-bonding in systems of hybrid materials including anilinium derivatives such as 4-Carboxyanilinium hydrogensulfate (Benali-Cherif, Direm *et al.*, 2007), 2-carboxyanilinium dihydrogenphosphate (Benali-Cherif, Allouche *et al.*, 2007) and *m*-Carboxyphenylanilinium bisulfate (Cherouana, *et al.*, 2003),

The asymmetric unit of (I) contains a monoprotonated *p*-hydroxyanilinium cation and bisulfate anion (Figure 1). Intra atomic bond distance and angles in the title compound shows the monprotonation of the organic entity and confirms the presence of the bisulfate (HSO_4^-) anion.

The crystal structure of the title compound is built up from intricate cation-cation, cation-anion and anion-anion hydrogen-bonds interaction in a three-dimensional network. Strong and moderate N—H···N, N—H···O, O—H···O hydrogen bonding ensure the cohesion of the crystal through the formation of three-dimensional hydrogen bond network and the strongest one are observed between cation and anion (O41—H···O4 2.695 (2) Å, (O1—H1···O41 2.642 (2) Å)). Principal hydrogen bonding values are listed in Table 1, and the interactions are illustrated in Fig. 2.

Experimental

Single crystals of the title compound are prepared by slow evaporation at room temperature of an aqueous solution of 4-hydroxyaniline acid $\text{C}_6\text{H}_7\text{NO}$ and sulfate acid (H_2SO_4).

Refinement

The OH and NH_3 H-atoms of the anion and cation entities were located in difference Fourier syntheses and refined as riding atoms with distances constraints of N—H = 0.89 Å and O—H = 0.82 Å [$U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{N}, \text{O})$]. Aromatic H atoms were located in difference Fourier syntheses and were allowed to ride on their parent C atoms with C—H = 0.93 Å and $U_{\text{iso}} = 1.2U_{\text{eq}}(\text{C})$.

Figures




Fig. 1. View of the title compound showing the atom-numbering scheme. Displacement ellipsoids are drawn at the 50% probability level. H bond is shown as dashed line. H atoms are represented as small spheres of arbitrary radii.





Fig. 2. Partial packing view showing the three dimensional hydrogen-bonding network.

3-Hydroxyanilinium hydrogensulfate

Crystal data



$M_r = 207.21$

Monoclinic, $P2_1$

Hall symbol: P 2yb

$a = 7.3142(3) \text{ \AA}$

$b = 5.8612(2) \text{ \AA}$

$c = 9.8969(2) \text{ \AA}$

$\beta = 98.829(2)^\circ$

$V = 419.25(2) \text{ \AA}^3$

$Z = 2$

$F_{000} = 216$

$D_x = 1.641 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

$\lambda = 0.71073 \text{ \AA}$

Cell parameters from 5862 reflections

$\theta = 2.1\text{--}30.0^\circ$

$\mu = 0.38 \text{ mm}^{-1}$

$T = 293(2) \text{ K}$

Prism, colourless

$0.15 \times 0.12 \times 0.10 \text{ mm}$

Data collection

Nonius KappaCCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 293(2) \text{ K}$

ω/θ scans

Absorption correction: none

5851 measured reflections

2241 independent reflections

2150 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.043$

$\theta_{\text{max}} = 30.0^\circ$

$\theta_{\text{min}} = 2.1^\circ$

$h = -8 \rightarrow 10$

$k = -7 \rightarrow 8$

$l = -13 \rightarrow 13$

Refinement

Refinement on F^2

Least-squares matrix: full

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$R[F^2 > 2\sigma(F^2)] = 0.030$	$w = 1/[\sigma^2(F_o^2) + (0.0598P)^2 + 0.051P]$
$wR(F^2) = 0.099$	where $P = (F_o^2 + 2F_c^2)/3$
$S = 1.14$	$(\Delta/\sigma)_{\max} = 0.001$
2241 reflections	$\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$
121 parameters	$\Delta\rho_{\min} = -0.45 \text{ e } \text{\AA}^{-3}$
1 restraint	Extinction correction: none
Primary atom site location: structure-invariant direct methods	Absolute structure: Flack (1983), with 1095 Friedel pairs
Secondary atom site location: difference Fourier map	Flack parameter: -0.02 (8)

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > 2\sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.2336 (2)	0.1606 (4)	0.28583 (15)	0.0237 (3)
C2	0.3143 (3)	0.3498 (4)	0.35270 (18)	0.0280 (4)
H2	0.3587	0.4675	0.3037	0.034*
C3	0.3287 (3)	0.3631 (3)	0.49435 (19)	0.0281 (4)
H3	0.3824	0.4902	0.5408	0.034*
C4	0.2629 (2)	0.1865 (4)	0.56539 (15)	0.0249 (3)
C5	0.1808 (3)	-0.0052 (4)	0.4976 (2)	0.0310 (4)
H5	0.1367	-0.1234	0.5463	0.037*
C6	0.1661 (3)	-0.0165 (4)	0.35594 (19)	0.0297 (4)
H6	0.1112	-0.1424	0.3088	0.036*
N1	0.2226 (2)	0.1419 (3)	0.13659 (15)	0.0271 (3)
H1A	0.2343	0.2798	0.1012	0.041*
H1B	0.1137	0.0826	0.1011	0.041*
H1C	0.3130	0.0520	0.1171	0.041*
O41	0.2772 (3)	0.1876 (3)	0.70515 (13)	0.0385 (3)
H41	0.2966	0.3181	0.7336	0.058*
O2	0.8476 (2)	0.0097 (3)	0.04947 (17)	0.0373 (4)
O1	0.8468 (2)	0.3880 (3)	0.13117 (15)	0.0333 (3)
H1	0.8085	0.4545	0.1941	0.050*
O3	0.5869 (2)	0.2501 (3)	-0.03018 (17)	0.0474 (5)
O4	0.6326 (2)	0.1115 (3)	0.20212 (15)	0.0334 (3)

supplementary materials

S1 0.71804 (5) 0.17750 (8) 0.08525 (3) 0.02299 (12)

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C1	0.0227 (6)	0.0291 (9)	0.0187 (6)	0.0027 (8)	0.0019 (5)	-0.0021 (7)
C2	0.0348 (10)	0.0259 (9)	0.0246 (8)	-0.0047 (8)	0.0085 (7)	0.0005 (7)
C3	0.0340 (9)	0.0245 (9)	0.0260 (8)	-0.0040 (7)	0.0056 (7)	-0.0035 (7)
C4	0.0294 (7)	0.0255 (8)	0.0202 (6)	0.0047 (9)	0.0046 (5)	-0.0006 (8)
C5	0.0390 (10)	0.0290 (10)	0.0253 (8)	-0.0072 (8)	0.0061 (7)	0.0017 (7)
C6	0.0341 (10)	0.0280 (10)	0.0261 (8)	-0.0056 (8)	0.0015 (7)	-0.0035 (7)
N1	0.0283 (6)	0.0325 (10)	0.0202 (6)	0.0031 (7)	0.0029 (5)	-0.0008 (6)
O41	0.0659 (9)	0.0314 (7)	0.0192 (5)	-0.0074 (9)	0.0092 (5)	-0.0022 (6)
O2	0.0310 (7)	0.0397 (9)	0.0432 (8)	0.0016 (7)	0.0120 (6)	-0.0167 (7)
O1	0.0339 (7)	0.0315 (8)	0.0366 (7)	-0.0064 (6)	0.0115 (6)	-0.0055 (6)
O3	0.0462 (9)	0.0469 (9)	0.0423 (8)	-0.0053 (8)	-0.0144 (7)	0.0167 (7)
O4	0.0407 (7)	0.0331 (8)	0.0303 (6)	-0.0025 (6)	0.0176 (6)	0.0004 (5)
S1	0.02315 (18)	0.0258 (2)	0.02035 (18)	0.00081 (18)	0.00441 (12)	0.00186 (16)

Geometric parameters (\AA , $^\circ$)

C1—C2	1.377 (3)	C6—H6	0.9300
C1—C6	1.382 (3)	N1—H1A	0.8900
C1—N1	1.4709 (19)	N1—H1B	0.8900
C2—C3	1.392 (3)	N1—H1C	0.8900
C2—H2	0.9300	O41—H41	0.8200
C3—C4	1.379 (3)	O2—S1	1.4467 (16)
C3—H3	0.9300	O1—S1	1.5759 (15)
C4—O41	1.3710 (18)	O1—H1	0.8200
C4—C5	1.396 (3)	O3—S1	1.4389 (15)
C5—C6	1.391 (3)	O4—S1	1.4491 (14)
C5—H5	0.9300		
C2—C1—C6	121.59 (15)	C1—C6—H6	120.3
C2—C1—N1	119.74 (18)	C5—C6—H6	120.3
C6—C1—N1	118.66 (18)	C1—N1—H1A	109.5
C1—C2—C3	119.31 (17)	C1—N1—H1B	109.5
C1—C2—H2	120.3	H1A—N1—H1B	109.5
C3—C2—H2	120.3	C1—N1—H1C	109.5
C4—C3—C2	119.56 (17)	H1A—N1—H1C	109.5
C4—C3—H3	120.2	H1B—N1—H1C	109.5
C2—C3—H3	120.2	C4—O41—H41	109.5
O41—C4—C3	122.2 (2)	S1—O1—H1	109.5
O41—C4—C5	116.7 (2)	O3—S1—O2	112.88 (11)
C3—C4—C5	121.13 (16)	O3—S1—O4	113.33 (10)
C6—C5—C4	118.94 (19)	O2—S1—O4	113.12 (10)
C6—C5—H5	120.5	O3—S1—O1	107.06 (10)
C4—C5—H5	120.5	O2—S1—O1	102.60 (9)
C1—C6—C5	119.47 (18)	O4—S1—O1	106.89 (9)

Hydrogen-bond geometry (Å, °)

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1A···O2 ⁱ	0.89	2.03	2.830 (2)	149
N1—H1B···O2 ⁱⁱ	0.89	1.98	2.853 (2)	166
N1—H1C···O3 ⁱⁱⁱ	0.89	2.14	2.962 (2)	152
N1—H1C···O4	0.89	2.39	2.975 (2)	124
O41—H41···O4 ^{iv}	0.82	1.88	2.695 (2)	174
O1—H1···O41 ^{iv}	0.82	1.86	2.642 (2)	160

Symmetry codes: (i) $-x+1, y+1/2, -z$; (ii) $x-1, y, z$; (iii) $-x+1, y-1/2, -z$; (iv) $-x+1, y+1/2, -z+1$.

Fig. 1




Fig. 2

